



THE WEIZMANN SCIENCE PRESS OF ISRAEL

National Council for Research and Development

ISRAEL Journal of BOTANY

Volume 23
1974



סד ויצמן לפרסומים במדעי הטבע ובטכנולוגיה, ירושלים

THE WEIZMANN SCIENCE PRESS OF ISRAEL

Publishers of the following journals

ISRAEL JOURNAL OF BOTANY; ISRAEL JOURNAL OF CHEMISTRY;
ISRAEL JOURNAL OF EARTH-SCIENCES; ISRAEL JOURNAL OF MATHEMATICS;
ISRAEL JOURNAL OF TECHNOLOGY; ISRAEL JOURNAL OF ZOOLOGY;
JOURNAL D'ANALYSE MATHÉMATIQUE

לדת מדע לנוער המתבגר (SCIENCE FOR YOUTH)

צע עתון מדעי לכל (SCIENCE)

Supporting Institutions

THE NATIONAL COUNCIL FOR RESEARCH AND DEVELOPMENT;
BAR-ILAN UNIVERSITY; BEN GURION UNIVERSITY OF THE NEGEV;
TECHNION—ISRAEL INSTITUTE OF TECHNOLOGY; TEL AVIV UNIVERSITY;
THE HEBREW UNIVERSITY OF JERUSALEM; THE WEIZMANN INSTITUTE OF SCIENCE

GENERAL EDITORIAL BOARD

D. ABIR
I. ARNON
A. DVORETZKY
H. EYAL-GILADI
H. HANANI
M. JAMMER
J. JORTNER
E. KATCHALSKI-KATZIR
N. SHARON
G. STEIN
Y. WEILER

Executive Editor

L. LESTER

Assistant Editor

ESME GORDON

ISRAEL JOURNAL OF BOTANY

Formerly Bulletin of the Research
Council of Israel, Section D

Editorial Board

LEONORA REINHOLD *Editor-in-chief*
NAOMI FEINBRUN
D. ATSMON
J. GALIL
N. LANDAU
G. ORSHAN
A. E. RICHMOND
Y. VAADIA
I. WAHL
Coordinator
U. PLITMANN

Subscriptions are to be addressed to the Weizmann Science Press of Israel, P.O.B. 801, Jerusalem 91000, Israel. In Europe, orders may be sent to Wm. Dawson & Sons Ltd., Cannon House, Macklin Street, London, W.C.2., or to booksellers. Subscription fees per volume, \$ 22.00, and beginning with Vol. 24, \$ 25.00.

Copyright © 1975 by THE WEIZMANN SCIENCE PRESS OF ISRAEL

Printed in Israel
Mercaz Press, Jerusalem

ISRAEL JOURNAL OF BOTANY

VOLUME 23, 1974

TABLE OF CONTENTS

Numbers 1-2

ANATOMY AND PHYSIOLOGY

- The problem of indehiscence of pistachio (*Pistacia vera* L.) fruit
A. Nevo, E. Werker and R. Ben-Sasson 1
- Leaf water potential and stomatal activity in sorghum as influenced by soil moisture stress
A. Blum and C. Y. Sullivan 14

GEOBOTANY

- Ecological studies on the vegetation of the Upper Galilee, Israel. I. A vegetation map of the Bar'am-Malkiyya area
A. Rabinovich-Vin and G. Orshan 20
- Cartographical methods and mapping with a computer
G. Halevy 38

GENETICS

- Non-brittle types in a wild population of *Triticum dicoccoides* Körn in Israel
A. Kamm 43

MYCOLOGY

- The Amanitaceae fungi of Israel
Z. Avizohar-Hershenzon and N. Binyamini 59

BIOLOGICAL FLORA OF ISRAEL

5. The subterranean clovers of *Trifolium* subsect. *calycomorphum* Katzn. *Trifolium subterraneum* L. (sensu latu)
J. Katznelson 69

BOOK REVIEWS

109

Number 3

ECOLOGY

- Ecological studies on the vegetation of the Upper Galilee, Israel. II. Factors determining the absence of batha and garigue components on Middle Eocene strata
A. Rabinovich-Vin and G. Orshan 111
- Effects of gazelles and seed beetles (Bruchidae) on germination and establishment of *Acacia* species
G. Halevy 120

ANATOMY AND PHYSIOLOGY

- Water potentials of stressed pine seedlings under controlled climatic conditions
D. Heth 127
- Heterocarpy and its ontogeny in *Aellenia austrani* (Post) Zoh. Light- and electron-microscope study
E. Werker and T. Many 132
- Ultrastructure of plastids of parasitic higher plants. V. Influence of light on *Cuscuta* plastids
G. Laudi, P. Medeghini Bonatti and G. Fricano 145

MYCOLOGY

- Pezizales of Israel. II. Pezizaceae
Z. Avizohar-Hershenzon and H. Nemlich 151

FLORISTICS

- Lathyrus sphaericus* Retz., a species new to Israel
J. Mattatia 164

SYSTEMATICS

The annual *Calendula* species: taxonomy and relationships

C. Heyn, O. Dagan and B. Nachman 16

ANATOMY AND PHYSIOLOGY

Some ecological aspects of distribution and germination of *Pancratium maritimum* L.

A. Keren and M. Evenari 20

Structure and organization of shoot apex of *Allium sativum* L.

I. L. Kothari and J. J. Shah 21

Intrusive growth in the phloem fibres of *Erythrina indica* and *Pongamia glabra*

A. K. M. Ghouse and D. Sabir 22

GEOBOTANY

Notes on the vegetation near Suez and Fayid (Egypt)

A. Danin 22

MYCOLOGY

Fleshy fungi of north and central Israel. I

N. Binyamini 23

BOOK REVIEWS

25

AUTHOR INDEX

Avizohar-Hershenzon, Z.,
59, 151

Ben-Sasson, R., 1
Binyamini, N., 59, 237
Blum, A., 14

Cohen, D., 254

Dagan, O., 169
Danin, A., 226

Evenari, M., 202

Fricano, G., 145

Ghouse, A. K. M., 223

Halevy, G., 38, 120
Heth, D., 127
Heyn, C., 169, 252

Kamm, A., 43
Katznelson, J., 69
Keren, A., 202
Kothari, I. L., 216

Laudi, G., 145

Many, T., 132
Mattatia, J., 164
Mayer, A. M., 255
Medeghini Bonatti, P., 145

Nachman, B., 169
Nemlich, H., 151
Nevo, A., 1
Noy-Meir, I., 252

Orshan, G., 20, 111

Rabinovich-Vin, A., 20, 111
Raven, P. H., 254

Sabir, D., 223
Shah, J. J., 216
Sullivan, C. Y., 14

Werker, E., 1, 132

SUBJECT INDEX

- Abies cephalonica*, 92
- abscission tissue, pistachio fruit, 1-13
- ascorbic acid, garlic shoot apex, 221
- Acacia*
- gerrardii* ssp. *negevensis*, 122, 123
- raddiana* 121, 123, 230, 233
- tortilis*, 122, 123, 233, 234
- Acacia* spp.,
- distribution in Negev and Sinai, computer and mapping, 38-42
- effect of gazelles and seed beetles, 120-126
- adaptation, *Acacia* spp., 120-126
- Aellenia autrani*, heterocarpy, 132-144
- Aethionema* spp., 142
- agronomy, subterranean clovers, 99-102
- Allium sativum*, shoot apex, 216-222
- Amanita*
- aspera*, 64-65
- boudieri*, 60
- echinocephala*, 60, 61
- eliae*, 65, 66
- inaurata*, 65, 67
- ovoidea*, 62
- pantherina*, 65, 66
- phalloides*, 62
- plumbea*, 67
- rubens*, 64
- rubescens*, 64
- vaginata* f. *vaginata*, 66-67
- verna*, 61, 63
- vittadini*, 60-62
- Amanitaceae, 245
- Amanitaria* Section, 66
- Amanitina* Section, 62-64
- Amanitopsis* Section, 66-67
- Amanitopsis strangulata*, 65, 67
- Amidella* Section, 62
- Amplariella* Section, 64-65
- Amygdalus*
- communis*, 27, 32
- korschinskii*, 27, 29, 31, 32
- ANABASETUM, 231, 232
- Anabasis articulata*, 228, 232, 233
- Anabasis articulata-Farsetia aegyptiaca* association, 229, 231
- Anagyris foetida*, 27, 30, 32
- Anastatica hierochuntica*, 233
- anatomy, heterocarpy in *Aellenia autrani*, 132-44
- apical elongation, phloem fibers, 223-225
- Artemisia monosperma*, 228, 230, 231, 232, 234
- Artemisia monospermum-Panicum turgidum* association, 228, 229
- Asparagus*
- aphyllus*, 27, 29, 30, 32
- stipularis*, 228, 233
- Aspidella* section, 60-62
- Avena*
- alba*, 30, 32
- sterilis*, 30, 32, 33
- Ballota*
- saxatilis*, 28
- undulata*, 28, 31, 32, 33
- barley, Mt. Tabor, 55-57
- batha, absence on Middle Eocene strata, Upper Galilee, 111-119
- batha communities, Upper Galilee, Bar'am-Malkiyya, 31-34
- biological flora, Israel, *Trifolium* spp., 69-108
- biotic factors, subterranean clovers, 91-96
- Bolbitiaceae, 248
- breeding
- Calendula* species, 169-201
- subterranean clovers, 169-201
- bruchid, pest of subterranean clover, 96
- Bruchidae, effect on germination and establishment of *Acacia* spp., 120-126
- Bruchidius*
- albosparsus*, 121
- spadiceus*, 124
- Bryonia*
- multiflora*, 27
- syriaca*, 27
- calcium chloride, effect on *Sarcopoterium spinosum*, 117
- Calendula*, spp., taxonomy and relationships, 169-201
- Calendula*
- aegyptiaca*, 170, 171, 173, 175, 180, 185, 191

- alata*, 173, 188–191
algeriensis, 174, 176, 178
arvensis, 169–175, 178, 180–191, 197–199
 aegyptiaca form, 185, 189, 197
 alata form, 188–191, 197
 persica form, 185, 197
 sancta form, 185–188, 189
 sinaica form, 185, 197
asterias, 176–178
bicolor, 170, 171
ceratosperma, 193
lanzae, 175
malacitana, 171
micrantha, 171
officinalis, 175
pachysperma, 171, 174, 175, 178, 193, 194–196, 198, 199
palaestina, 171, 174, 175, 178, 191–196, 198, 199
persica, 171, 173, 185
platycarpa, 174, 180
repanda, 193
sancta, 169, 170, 173, 185–188
sinaica, 171, 185
stellata, 174, 176–178, 197–198, 199
triptocarpa, 174, 175, 178–180, 197–198, 199
Calligonum comosum, 232
Calycotome villosa, 28, 30–32, 35, 118, 152
Capparis spinosa, 28, 30, 32
Capra ibex nubiana, 120–126
 carbohydrates, garlic shoot apex, 221
Carlina
 corymbosa, 92
 involucrata, 33
 cartographical methods, computer, 38–42
Caryedon serratus, 122
Centaurea aegyptiaca, 229, 233
Ceratonia siliqua, 27, 28, 29
Cercis siliguastrum, 27, 28, 29
 chemotaxonomy, subterranean clovers, 76
Chenopodium spp., 142
 chlorophyll content, *Cuscuta australis*, 145–150
 chloroplasts, *Cuscuta australis*, 145–150
 chromosome counts, *Calendula* species, 169–201
Cistus, 26, 152
 creticus, 30
 salviifolius, 28
Cistus spp.-*Calycotome villosa* garigue, Upper Galilee, Bar'am-Malkiyya, Israel, 34
Clematis
 cirrhusa, 27
 flammula, 27
 climate
 Fayid and Suez, 226–227
 Upper Galilee
 water potential and pine seedlings, 127–131
 climate and distribution, subterranean clover, 87–89
Clitocybe subalutacea, 238
 clovers, subterranean, biological flora, Israel, 69–108
 comb. nov.
 Peziza
 nivalis, 162
 succosella, 156–157
 computer, cartographical methods and mapping, 38–42
Conocybe ambigua, 248
Convolvulus lanatus, 228, 231, 232, 234
Convolvulus lanatus-Heliotropium digynum association, 229, 231
 Coprinaceae, 245–248
Coprinus
 atramentarius, 245–246
 domesticus, 246
 xanthothrix, 246
Cornulaca monacantha, 228, 232
 Cortinariaceae, 248–250
Crataegus aronia, 27, 29, 30, 32
 Crepidotaceae, 250–251
Crepidotus
 luteolus, 250–251
 pubescens, 250–251
 sphaerosporus, 250
Crotolaria
 aegyptiaca, 233
Cupressus
 arizonica, 152
 macrocarpa, 152
Cuscuta australis plastids, ultrastructure, 145–150
Cynodon dactylon, 92
 cytogenetics
 Calendula species, 169–201
 Trifolium subsect. *Calycomorphum*, 72–76

- cytology
Calendula species, 169–201
 garlic shoot apex, 216–222
- Dactylis glomerata*, 30, 32, 33
- development
 heterocarpy in *Aellenia autrani*, 132–144
 shoot apex, garlic, 216, 222
 subterranean clovers, 76–87
- diaspores, *Aellenia autrani*, 132–144
- Dimorphotheca*, see *Calendula*
- diseases, subterranean clover, 95–96
- distribution
Acacia spp., Negev and Sinai, computer mapping, 38–42
 Amanitaceae, Israel, 59–68
Calendula species, 169–201
 Pezizaceae, Israel, 151–168
Trifolium subsect. *Calycomorphum*, 71–72, 87–96
- DNA, shoot apex, garlic, 220, 221
- domestication, emmer, 54–57
- dwarf shrub communities, subterranean clover, 92
- Echinops*, 92
gaillardotii, 33
spinosus, 228, 230, 232, 234
- Echinops spinosus-Ephedra alata* association 232
- ecological studies, Upper Galilee, 20–37
- ecology
 seed germination and dispersal, *Panocratium maritimum*, 202–215
 wild emmer, 54–57
- ecology and distribution, subterranean clovers, 87–90
- edaphic factors and distribution, subterranean clovers, 89–90
- Egypt, vegetation, near Suez and Fayid, 226–236
- electron microscope, heterocarpy in *Aellenia autrani*, 132–144
- embryo, *Aellenia autrani*, 137–139
- emmer, cultivated and wild, Israel, 43–58
- endocarp, pistachio fruit, indehiscence, 1–13
- Ephedra*
alata, 232
campylopoda, 27, 29, 32
Erodium hirtum, 228, 233
Erysiphe polygonii, 96
Erythrina indica, intrusive growth of phloem fibres, 223–225
 establishment, *Acacia* spp. effect of gazelles and seed beetles, 120–126
 estrogenic activity, subterranean clovers, 96–99
Euphorbia hierosolymitana, 28, 31, 33
- Fagonia*
arabica, 228
bruguieri, 233
- FAGONIETUM ARABICAE, 229
- Farsetia aegyptiaca*, 228–232
- Fayid, vegetation, 226–236
- Fibigia clypeata*, 33
- Ficus*
pseudo-sycomorus, 233
retusa, 152
- flowering, subterranean clovers, 78–81
- Foeniculum vulgare*, 32
- forest associations, subterranean clovers, 92
- forest plant communities, Upper Galilee, Bar'am-Malkiyya, 26–31
- fruit indehiscence, pistachio, 1–13
- Fumana arabica*, 30
- fungal diseases, subterranean clover, 69
- fungi
 Amanitaceae, Israel, 59–68
 fleshy, North and Central Israel, 237–251
 fusiform initials, phloem fibers, 223–225
- Galactina*, see *Peziza*
- garigue, absence on middle Eocene strata, Upper Galilee, 111–119
- garigue communities
 subterranean clovers, 92
 Upper Galilee, Bar'am-Malkiyya area, 13–14
- garlic, shoot apex, 216–222
- Gazella dorcas*, 120–126
- gazelles, effect on germination and establishment of *Acacia* spp., 120–126
- genetics
Calendula species, 169–201
 emmer and tetraploid wheat, 43–58

- Trifolium* subsect. *Calycomorphum*, 72–76
- germination
- Acacia* spp., effects of gazelles and seed beetles, 120–126
- Pancreatium maritimum*, 202–215
- Sarcopoterium spinosum*, 115
- subterranean clovers, 83–84
- grazing and subterranean clovers, 91
- growth
- intrusive, phloem fibers, *Erythrina indica* and *Pongamia glabra*, 223–225
- shoot apex, garlic, 216–222
- subterranean clovers, 76–87
- Gymnarrhena micrantha*, 142
- Gymnocarpus decander*, 229, 232, 233
- Gymnocarpus decander-Stachys aegyptiaca* association, 229–233
- habitats
- Calendula* species, 169–201
- Pezizaceae, Israel, 151–168
- Helianthemum kahiricum*, 228, 232, 233
- Heliotropium digynum*, 228, 231, 234
- hemicyptophytic batha, subterranean clover, 92
- heterocarpy, *Aellenia autrani*, 132–144
- histochemistry, shoot apex, garlic, 220–221
- histology, shoot apex, garlic, 216–222
- Hohenbuehelia*
- geogenius*, 242
- longipes*, 241–242
- Hordeum*
- bulbosum*, 32, 33
- ithaburens*, 33
- spontaneum*, 33
- Hordeum bulbosum-Euphorbia hierosolymitana* association, 33, 34
- Majorana syriaca* variant, 34
- Smyrnanium connatum* variant, 34
- hybridization, effect on pistachio fruit indehiscence, 2–13
- Hyoscyamus muticus*, 228
- Hyparrhenia hirta*, 234
- ibex, 120–126
- indehiscence, pistachio fruit, 1–13
- Inocybe*
- fastigiata*, 249
- furfurea*, 249
- gramata*, 249–250
- jurana*, 248–249
- Iphiona mucronata*, 229, 233
- Iphiona mucronata-Gymnocarpus decander* association, 229
- isoflavones, occurrence and genetics, subterranean clover, 96–99
- karyotypes, *Calendula* spp., 169–201
- key
- annual *Calendula* spp., 174, 175
- Trifolium* subsect. *calycomorphum*, 69–70
- Kickxia aegyptiaca*, 229, 232
- Lasiurus hirsutus*, 229, 232, 233
- Lathyrus*
- inconspicuus*, 164, 166
- sphaericus*, new to Israel, 164–166
- Laurus nobilis*, 27
- leaf diffusion resistance, Sorghum, 14–19
- leaf water potential, sorghum, 14–19
- light
- effect on plastids, *Cuscuta australis*, 145–150
- seed germination, *Pancreatium maritimum*, 206, 210, 213
- Limonium pruinosum*, 229, 234
- Lonicera etrusca*, 27
- Lycium shawii*, 229, 234
- Lyophyllum*
- aggregatum*, 241
- crassifolium*, 241
- Majorana syriaca*, 28, 30–34
- Malabala sekakul*, 33
- man and distribution of subterranean clovers, 91
- mapping, computer, 38–42
- maquis communities
- subterranean clovers, 92
- Upper Galilee, Israel, 26–31
- Marasmiellus albus-corticis*, 243–244
- Marasmius*
- candidus*, 243–244
- cohaerens*, 244
- Medicago polymorpha*, 32

- Melanoleuca*
excissa, 241
humilis, 238
schumacheri, 240
stridula, 240
- microfibrils, pistachio fruit endocarp, 7, 9
- Middle Eocene strata, absence of batha and garigue components, Upper Galilee, 111–119
- moisture stress, sorghum, 14–19
- morphology, emmer, 47–58
- mutation, wild and cultivated emmer, 56–57
- Mycena hiemalis*, 244
- Naucoria pellucida*, 250
- needle water potential, pine seedlings, 127–131
- Negev, *Acacia* spp., 120–126
- new records in Israel
Amanitaceae, 59–68
 fleshy fungi, 237–251
Pezizaceae, 151–163
- new to Israel
Lathyrus sphaericus, 164–166
- Noaea mucronata*, 233
- nom. nov., *Peziza moseri*, 157–158
- nucleic acids, shoot apex, garlic, 220–221
- nutritive value, subterranean clover, 96
- oak trees, fleshy fungi, 237–251
- obituary, Gideon Halevy, 167–168
- Ochrademus baccatus*, 234
- Oidium* spp., 96
- Olea europaea*, 17
- Ononis*
antiquorum, 29, 30, 33
natrix, 28, 30–33
- Ononis natrix-Sideritis perfoliata* association, 31–34
- ontogeny, shoot apex, garlic, 216–222
- Orchidaceae, 31
- origin, cultivated wheat, 57
- Oryzopsis*
holiciformis, 30, 32, 33
miliacea, 30, 234
- Osteospermum*, 196
- Osyris alba*, 28, 30, 32, 33
- Paliurus*, 92
- Pancratium matitinum*, seed germination and dispersal, 202–215
- Panicum turgidum-Convulvulus lanatus* association, 229
- Panicum turgidum-Farsetia aegyptiaca* association, 229
- Panus conchatus*, 238
- parasitic higher plants, ultrastructure of plasmids, 145–150
- Paronychia desertorum*, 229
- perianth, *Aellenia autrani*, 133–135
- pericarp, *Aellenia autrani*, 135–136
- pests, subterranean clovers, 95–96
- Peziza*
brunneoatra, 153, 159, 161–162
cerea, 156
cervina, 153, 159, 161
fuliginea, 153–155
micropus, 153, 159, 160
moseri, 153, 155, 157–158
nivalis, 153, 162
proteana forma *sparassoides*, 153, 155, 158
repanda, 153, 159, 160
saniosa, 153, 155, 156
sepiatra, 153, 159, 161
succosella, 153, 155, 156–157
succosa, 153, 155, 165
tectoria, 156
varia, 153, 158–160
 forma *typica*, 159, 160
vesiculosa, 151, 153, 159, 161
violacea, 153, 155, 157
- Pezizaceae*, Israel, 151–163
- Pezizales*, Israel, 151–163
- Phagnalon*
barbeyanus, 234
rupestre, 28, 30, 32, 34
rupestris, 229, 233, 234
- Phillyrea media*, 27
- phloem fibers, intrusive growth, 223–225
- Phlomis viscosa*, 28, 30
- phytosociology, subterranean clovers, 91–93
- pine seedling, water stress, 127–131
- pine trees, fleshy fungi, 237–251
- Pinus*
brutia, water stress, 127–131
halepensis, 152
nigra, 92
radiata, water stress, 127–131

- taeda*, water stress, 127–131
- Pistacia*
- atlantica*, 27–28, 29, 31
- palaestina*, 27–32, 34
- vera*, fruit indehiscence, 1–13
- Pistacia atlantica*-*Amygdalus korschinskii* association, 29, 31, 36
- pistachio fruit, indehiscence, 1–13
- Pituranthos tortuosus*, 229, 232
- Pituranthos tortuosus*-*Panicum turgidum* association, 229, 232
- plastids, ultrastructure, *Cuscuta australis*, 145–150
- Pleurotus*
- cornucopiae*, 237–238
- longipes*, 241–242
- Pluteus*
- godeyi*, 245
- nanus*, 245
- POETO-TRIFOLIETUM SUBTERRANEI, 93
- Polyporaceae, 237
- Pongamia glabra*, phloem fibers, 223–225
- population structure, *Trifolium* subsect. *Calycomorphum*, 72–74
- Potentilla* spp., 92
- Prangos ferulacea*, 34
- Prasium majus*, 27
- Prunus ursina*, 26–31
- Psathyrella*
- candolleana*, 246
- conopilea* var. *subatrata*, 246
- fatua*, 248
- pannucoides*, 246–248
- spadiceo-grisea*, 248
- Psoralea betuminosa*, 33
- Pyrus syriaca*, 26–31
- QUERCETALIA ILICIS, 92
- QUERCETUM CALLIPRINI, 37
- Quercus*
- boissieri*, 26–29
- calliprinos*, 26–31, 34, 152
- ilex*, 92
- ithaburensis*, 92
- pubescens*, 92
- suber*, 92
- Quercus calliprinos*-*Pistacia palaestina* association, Upper Galilee, 20–37
- Bar'am variant, 26–27
- Pistacia atlantica* variant, 28–29
- Pyrus syriaca* variant, 26
- Quercus boissieri* variant, 26–28, 35
- Styrax officinalis* variant, 20–29
- Reaumuria hirtella*, 229
- Retama raetam*, 234
- Rhamnus*, 34
- alaternus*, 27, 28, 32
- palaestinus*, 27, 28, 30, 32
- punctatus*, 29, 30, 32
- Rhizobium* and subterranean clovers, 93–95
- Rhodophyllaceae*, 244, 245
- Rhodophyllus*
- asprellus*, 244
- cancrinus*, 245
- sericeus*, 244
- RNA, garlic shoot apex, 220–221
- Rubia tenuifolia*, 27, 29, 30
- Rubus sanctus*, 28
- Ruscus aculeatus*, 28
- Ruta chalapensis*, 32
- salinity, seed germination, *Pancratium maritimum*, 211–214
- Salsola*
- inermis*, 142
- tetrandra*, 229, 233–235
- SALSOLETUM TETRANDAE, 229, 223
- Salvia*
- dominica*, 30, 32
- triloba*, 20, 26
- sap pressure, pine seedlings, 127–131
- Sarcopoterium spinosi*, 30–31, 34, 35
- Sarcopoterium spinosum*, 26, 28, 30, 31, 36, 92, 111–119
- Sarcosphaera crassa*, 152–153, 155
- Scrophularia xanthoglossa*, 32, 234
- Sedum* spp., 34
- seed beetles, effect on germination and establishment of *Acacia* spp., 120–126
- seed dispersal, *Pancratium maritimum*, 202–215
- seed dormancy, subterranean clovers, 85–87
- seed germination
- Acacia* spp., effect of gazelles and seed beetles, 120–126

- Pancratium maritimum*, 202–215
 subterranean clovers, 83–84
 seed impermeability, subterranean clovers, 84–85
 seed size, pistachio fruit, 1–13
 seedset, subterranean clovers, 81–83
 shoot apex, *Allium sativum*, 216–222
Sideritis perfoliata, 28, 30–33
 Sinai, *Acacia* spp., 120–126
Smilax aspera, 27–30
Smyrniolum connatum, 35
 soil moisture and distribution, subterranean clover, 88–89
 soil moisture relationships, Middle Eocene terra rossa, Upper Galilee, 112–113
 soil moisture stress, 14–19
 soil water potential
 pine seedlings, 127–131
 sorghum leaf, 14–19
Sorghum bicolor, soil moisture stress, 14–19
Spartium junceum, 26, 27, 30, 31, 34
Stachys aegyptiaca, 229, 233
Stipa parviflora, 229, 233
Stipagrostis
 plumosa, 229, 231
 raddiana, 233
 stomatal activity, sorghum, 14–19
 stomatal pore width, sorghum leaf, 14–19
 stress, soil moisture, sorghum, 14–19
 structure, shoot apex, garlic, 216–222
Styrax officinalis, 27, 29, 30, 32
 subterranean clovers, biological flora, Israel, 69–108
 Suez (Egypt), vegetation, 226–236
 survival, *Sarcopoterium* seedlings, 114–115
- Tamus*
 communis, 27, 30
 orientalis, 27
Taverniera aegyptiaca, 233, 234
 taxonomy
 Calendula species, 169–201
 Pezizaceae, Israel, 151–168
 Trifolium subsect. *calcycomorphum*, 69–70
 temperature, seed germination, *Pancratium maritimum*, 205–206, 212–213
 terra rossa, Middle Eocene strata, Upper Galilee, 111–119
 testa, *Aellenia autrani*, 137
Teucrium polium, 30, 32–33, 34
 transpiration, sorghum leaf, 14–19
Tricholoma
 crassifolium, 241
 saponaceum, 238
 strictipes, 240
 Tricholomataceae, 238–244
Trifolium
 brachycalycinum, 69–108
 var. *brachycalycinum*, 70
 var. *flagelliforma*, 70
 var. *graecum*, 70
 var. *oxaloides*, 70
 campestre, 30, 32
 clusii, 32
 fragiferum, 92
 israeliticum, 69–108
 purpureum, 30, 32
 resupinatum, 30, 32
 subterraneum, biological flora, Israel, 69–108
 yanninicum, 69–108
Trifolium subsect. *calcycomorphum*, biological flora, Israel, 69–108
Trigonella lilacina
Triticum
 dicoccoides, Mt. Gilboa, Israel, 43–58
 dicoccum, 55
 durum, 54, 56
 triccoccum, 56
Tubaria
 autochthona, 250
 furfuracea, 250
 pallidospora, 250
- ultrastructure
 Cuscuta australis plastids, 145–150
 heterocarpy in *Aellenia autrani*, 132–144
 Upper Galilee, vegetation, 20–37, 111–119
Uromyces sp., 96
- variation, *calendula* species, 169–201
Varthemia, 34
 iphionoides, 28, 30
 montana, 233
Varthemietum iphionoides community, 34
 vegetation
 Suez and Fayid, Egypt, 226–236

Upper Galilee, 20-37, 111-119
vegetation map, Upper Galilee, 20-37
Viability, *Pancratium maritimum* seeds, 202-215
Vicia peregrina, 30-32
viral diseases, subterranean clover, 96

water potential

pine seedlings, 127-131
sorghum leaf, 14-19
water relations, sorghum plants, 14-19
water stress, pine seedlings, 127-131

wheat, tetraploid, 43-58

xylem pressure potential, pine seedlings, 127-128

Zilla spinosa, 229

Zygophyllum

coccineum, 232, 233
decumbens, 234
dumosum, 234